



Early Journal Content on JSTOR, Free to Anyone in the World

This article is one of nearly 500,000 scholarly works digitized and made freely available to everyone in the world by JSTOR.

Known as the Early Journal Content, this set of works include research articles, news, letters, and other writings published in more than 200 of the oldest leading academic journals. The works date from the mid-seventeenth to the early twentieth centuries.

We encourage people to read and share the Early Journal Content openly and to tell others that this resource exists. People may post this content online or redistribute in any way for non-commercial purposes.

Read more about Early Journal Content at <http://about.jstor.org/participate-jstor/individuals/early-journal-content>.

JSTOR is a digital library of academic journals, books, and primary source objects. JSTOR helps people discover, use, and build upon a wide range of content through a powerful research and teaching platform, and preserves this content for future generations. JSTOR is part of ITHAKA, a not-for-profit organization that also includes Ithaka S+R and Portico. For more information about JSTOR, please contact support@jstor.org.

which these specimens were gathered, was surrounded by others, the culms of these surrounding ones interlacing those of the plant exhibited, but only this one plant was infected. He did not count the number of culms, but felt safe in saying there were over fifty. In walking through this field among many hundreds of plants of this *Panicum*, he saw only one other plant, which in like manner was infested. This had one perfect panicle only among the numerous infested ones—the interlacing branches of surrounding plants of the same species being free, as in the other instance. It was scarcely credible that sporidia of the *Ustilago*, floating through the atmosphere, settled on fifty separate culms of one plant, and not one on the culms of adjacent plants which were growing in and among them. Again, the leaves of the *Panicum* have a large spathaceous sheath, two or three inches long. The *Ustilago* attacked the panicle while closely swathed in this sheath, and fully perfected its growth entirely therein. He had indeed to unfold the sheath in order to detect the mass of “smut” to which the embryonic panicle was reduced, in order to detect its presence. Only the peculiar appearance of the grassy tuft having no inflorescence as in the case of its neighbors, drew attention to the plant in the first instance. If it seemed incredible that fifty culms interlocked with as many from other plants, should each receive a germinating spore alone, it was still more incredible that the spores should have found their way from the outside to the interior of these tightly twisted sheaths.

These observations did not prove that the sporidia entered the plant by the roots, and made their way in some incomprehensible manner through the structure to the inflorescence; but they did render the external-entrance hypothesis doubtful, and, in connection with Queckett's experiments, are possibly of some worth.

Dr. LEIDY made some remarks on Mr. Meehan's communication, showing that the tendency of modern observations rather favored the view that the entrance of the sporidia of microscopic fungi was from the outside.

Sexual Characters in Cephalotaxus.—MR. MEEHAN exhibited some fruit of *Cephalotaxus Fortunei*, a Chinese tree, this plant growing on the grounds of P. J. Berckmans, at Augusta, Georgia. This tree had for many years produced male flowers only. During 1882, it produced abundance of fruit. It showed that the genus was not truly dioecious, and further it afforded an illustration now not uncommon, that trees a long time of one sex only, would sometimes change to another. Sex is not an invariable characteristic in an individual tree.

A New Infusorian belonging to the Genus Pyxicola.—Prof. LEIDY exhibited drawings of an infusorian, a species of *Pyxicola*, which appeared to be different from those previously described.